



## Soil Organic Matter and its Importance for Water Management

This Deltafact highlights the role of soil organic matter in retaining water in the soil (sponge effect) and the options to influence its quantity and what this implies for water management.

1. INTRODUCTION
2. RELATED TOPICS AND DELTAFACTS
3. STRATEGY
4. SCHEMATIC REPRESENTATION
5. INTERACTIONS
6. COSTS AND BENEFITS
7. PRECONDITIONS AND LOCATIONS
8. GOVERNANCE
9. FIELD EXPERIENCES
10. KNOWLEDGE GAPS
11. SOURCES & LINKS
12. OVERVIEW CURRENT INITIATIVES AND RESEARCH
13. DISCLAIMER

### 1. Introduction

This Deltafact depicts the background of functions of soil organic matter, options to influence its quantity in the soil on agricultural plots and what implications are for water management of an increase of soil organic matter content, while presenting a scientific basis for this.

Stakeholders from a number of sectors have expressed concern about the decline in organic matter quantity in Dutch arable land. Time series analyses ([van den Akker](#),

2012 and Eurofin Agro, 2017), however, show neither downward nor upward trends on a national level. Locally there may be an ascending or descending soil organic matter quantity, for example when grassland is converted into arable land, or vice versa ([Smit et al., 2007](#)). However, the composition and quality of soil organic matter is changing structurally ([Eurofins, 2017](#)).

Sufficient soil organic matter is fundamentally important to availability of water and nutrients, trafficability, carbon sequestration, resilience against diseases and plagues and crop production. Maintaining and, where needed, increasing soil organic matter content serves to meet challenges that intensively used agricultural lands face, like dealing with extreme precipitation and drought, both occurring more frequently due to climate change.

That the organic matter content of agricultural fields should increase, has also landed as an issue in politics these days. The Dutch Minister Schouten (Agriculture and Nature) has recently stated in her vision about circular agriculture: "*A soil containing much organic matter, is better equipped to absorb water and is more resistant against drought. Such a soil can also retain more nitrogen and minerals, offers a richer soil life and contributes to healthy crops.*" ([Schouten, 2018., p.22](#)). The Soil Strategy, 2016 (in Dutch) of the [Soil Technical Commission \(TBC\)](#) of the Dutch Government indicates specific strategies for managing organic matter on agricultural fields. The draft Climate Accord ([Klimaatberaad, 2018](#)) refers to the significance of more organic matter in the soil.

## 2. Related topics and Deltafacts

**Topics:** Drought damage, Wet damage, Soil life, Soil structure, Soil compaction, Water shortage and freshwater supply, Water quality, Water storage

**Deltafacts:** [Bodem als buffer](#), [Onderwaterdrains](#), [Bodemvocht gestuurd beregenen](#), [Droogte stuurt functies](#),

## 3. Strategy

Soil organic matter receives attention nowadays as an option of improving the management of water at plot level and, at higher scales, of buffering effects of climate change on the water system. A healthy soil can provide various ecosystem services, linked to Retention, Storage and Supply ([RIVM, 2014; Bodem als buffer](#)). This notion may induce farmers, on behalf of a proper water management, to

maintain the soil organic matter quantity or to even increase this, by applying adaptive cultivation systems and by supplying extra organic material. In this context, it is always important for the water manager, to question what these measures will mean on an area level to the water availability in dry periods, to the surface run-off in wet periods and to the water quality.

#### 4. Schematic representation

Figure 1 shows organic matter content in Dutch top soils and helps identifying areas that are vulnerable with regard to their water availability. Figure 2 gives the many soil interactions related to organic matter.




Figure 1. Organic matter content at 15 cm depth in arable soils in the Netherlands ([Van den Berg et al., 2017](#)) and the profile of a podzol in which the dark upper layer indicates presence of organic matter.




Figure 2. Schematic representation of soil interactions related to organic matter; blue lines: effect of soil life; green arrows: soil physical effects; red lines: soil chemical effects; dark lines: effects of soil processes. Dash lines: correlation with organic matter. Source: [Reubens et al., 2010](#)

## 5. Interactions

### Organic matter, nutrient availability and soil structure

The influence of the quantity of soil organic matter on the nutrient balance is complex. An increase of the organic matter content leads to a stronger soil bonding of nutrients due to a higher cation-exchange capacity (CEC). When organic matter is decomposing, nutrients become available, so less fertiliser will suffice. However, in case nutrients are released in a period in which the crop does not require them, a higher organic matter content may lead to larger nutrient losses to ground- and surface water. Generally, clay soils have a good natural soil fertility. Sandy soils

strongly depend on organic matter for nutrient supply. In these soils availability of nutrients is largely determined by interactions between soil life and soil organic matter.

There is a clear correlation between organic matter content and soil structure ([Faber et al., 2011](#)). The more organic matter present, the better soil structure will be. A good soil structure is essential for the carrying capacity and infiltration capacity, it limits the sensitivity for sealing and the chance of soil compaction, it makes the soil less sensitive for soil diseases and gives a higher crop yield.

### Organic matter and water availability

The influence of the quantity of organic matter on water availability may be exerted directly or indirectly.

#### *Direct effect on water retention*

A direct effect is the influence of organic matter on the water retention. The water availability of the soil is calculated, by multiplying the quantity of available water between field capacity at  $pF = 2$  and wilting point at  $pF=4.2$ , with the thickness of the root zone  $\Delta z$ , like indicated in figure 3.




Figure 3. Water retention characteristic indicating the relationship between the pressure head ( $pF$  value) and the corresponding volumetric moisture content ( $\Theta$ ).

Derived from information about sandy soils in the Staring series ([Wösten et al., 2001](#)), the relation is calculated between the organic matter content and moisture content when saturated, at pF 2 (field capacity) and at pF 4.2 (wilting point). Figure 4 shows the result of calculations for a sandy soil and indicates that moisture content at all 3 pressure heads increases with increasing organic matter contents.




Figure 4. Curve of the volumetric moisture content at saturation, at pF 2 (field capacity) and at 4.2 (wilting point) with organic matter content

Figure 5 shows, on the basis of the above calculations, the curve of the soil water availability in relation to the organic matter content of a sandy soil with two different loam contents. The water availability here is the quantity of water which the plant can take up from the soil between field capacity (pF 2) and wilting point (pF 4.2).

Taking figure 5 as the starting point, figure 6 gives the increase of the water availability with increasing organic matter contents.

Since the sandy soils from the Staringgreeks ([Wösten et al., 2001](#)) contain minimally 1% organic matter, water availability itself (figure 5) and its increase (figure 6) are estimations. In case organic matter contents are below 1%, these are shown, for this reason, as dotted lines.




Figure 5. Curve of the water availability and the organic matter content.




Figure 6. Increase of water availability (mm) in a layer of 10 cm with an increase of organic matter of 1%.

#### Indirect effect on water retention

An indirect effect is that organic matter lowers the bulk density of soils and therefore the resistance against penetration by plant roots. Plants on soils with higher organic

matter contents may not only take up more water from a certain root zone, but this root zone can also become thicker, making again more water available.

**The most important conclusions in relation to organic matter content and water availability are:**

- An increase of organic matter content leads to a direct and indirect increase in water availability. The indirect effect has not been quantified.
- Poor sandy soils with 0,5 up to 1 % organic matter will render, with an increase of 1 % organic matter, an increase of 3 – 4 mm available water (figure 6) in a root zone with a thickness of 20 cm. This roughly equals one day extra transpiration.
- In the trajectory from 1 to 3 % organic matter, an increase of 1 % organic matter leads to an increase of 2 -3 mm available water (figure 6) in a root zone of 20 cm this equals less than one day extra transpiration.
- In the trajectory from > 3 % organic matter, an increase of 1 % organic matter leads to an increase of 1 mm available water in a root zone of 20 cm (sand).
- In a dry summer, in which the precipitation deficit continuously increases, extra water availability will have little effect. However, in a moderately dry summer with a regular shower, extra water availability of 3-4 mm can be used multiple times and it could lead to postponing a sprinkler irrigation application of 20 – 25 mm ([Bodemvocht gestuurd beregenen](#)) or to not apply this at all.
- Increasing organic matter content will, in the Netherlands, have the greatest impact on water availability in the green- and yellow coloured sandy soils in the mid- and eastern part of North Brabant and in the Achterhoek and Twente (Figure 1). The nature areas (for example the Veluwe) and coastal and riverine clay soils are not relevant in this. The agricultural fields on clay already have a large water retention capacity, regardless its organic matter content ([Van den Berg et al., 2017](#)).
- The more organic matter is present, the better the soil structure will be, thereby giving the soil a higher infiltration capacity with less ground level run-off (indirect positive effect on the water retention and possibly reduction of peak discharges ([Schipper et al., 2015](#)). Also the composition and the ratio between dynamic and stable organic matter can have an influence on the water retention ([Eurofins, 2017](#)). No field studies are known, that quantify these indirect effects for Dutch soils, yet there are indications from field research elsewhere ([Williams et al., 2017](#)), that point to a positive effect.

- A good soil structure makes it possible that plants root well, which means increasing water availability for the crops. Based on an average of 25 % available water (figure 5), a root zone of 20 cm renders 50 mm water. With a root zone of 30 cm this is 75 mm. This means an increase of 50%.

### **Ways to increase organic matter content**

First of all, it is important to quantify what an increase of 1 - 5 % organic matter means in terms of the yearly supply of organic matter. It is important hereby, to make a distinction between two forms of organic matter: old organic matter ("humus") that degrades at a rate of 2% under Dutch circumstances, and fresh organic matter of which 40 % is converted into humus within one year (Figure 7).

$$H = H_E + (H_0 - H_E) \exp(-\alpha t)$$

$H_E$  is evenwichts waarde van humus,

$$H_E = \frac{pI}{\alpha}, p = \text{omzettings fractie van verse organische stof } (0.4 \text{ jr}^{-1})$$

$$\alpha = \text{afbraaksnelheid humus } (0.02 \text{ jr}^{-1})$$

$$I = \text{Jaarlijkse aanvoer van verse OS } (\text{kg ha}^{-1} \text{ jr}^{-1})$$

$$H_0 = \text{initieel humus } (\text{kg ha}^{-1})$$

Figure 7. Calculation method for the volume of supplying humus (in kg per hectare)

Fresh organic matter is superficially supplied and subsequently ploughed through. Conversions will then take place in the 20-25 cm thick root zone. Such a root zone corresponds with ca  $3 \cdot 10^6$  kg soil and 1% of this is  $3 \cdot 10^4$  kg ha $^{-1}$ . The yearly supply would then amount to  $0.02 \cdot 3 \cdot 10^4 / 0.4 = 1,500$  kg organic matter per hectare, which is quite a significant volume. This is why it is more practical to follow a long-term strategy, instead of aiming to reach this 1% increase in 1 year. The strategy would then be to increase the organic matter content by 1% gradually, over several years.

The soil organic matter content, in general, can be raised in the following 3 ways ([SmartSoil project, 2015](#)):

1. Advancement of input in the soil of crop- and root residues
2. Increase of the quantity of soil organic matter by supplying manure and compost (both from external sources as from the farm itself)
3. Reduction of de-composting losses by limiting disturbance of the soil

The following cultivation measures have a positive effect on conserving and increasing of the soil organic matter content:

- **Crop rotation**

Cultivation of crops with a long growing season combined with the cultivation of leguminous plants that improve the quality of organic matter.

- **Crop residues**

Crop residues are those materials that are left on the field after the crop has been harvested. This includes stems, stubbles, leaves, roots and chaff. Insert these crop residues maximally into the soil.

- **Supply of manure and compost**

Supply of manure and compost effectively enhances the organic matter content, because it decomposes less quickly than fresh crop- and root residues. Supplying manure and compost will also often decrease the necessity to apply fertiliser.

- **Soil covering and catchment crops**

Prevention of fallowing in winter, by cultivating crops year-round, enhances the insertion of organic matter and decreases soil erosion and leaching of nutrients.

- **Conservation agriculture**

Conservation agriculture consists of minimizing soil tillage and having permanent soil coverage and diversity in crops. Minimal- and eventually no soil tillage, by a transition from frequent ploughing to forms of non-inversion tillage, diminishes the decomposition of organic matter.

The five measures are summarised in Table 1. Both the prevention of fallowing and the reduction of tillage take time before it results in an obvious increase of organic matter content.

<b>Measure</b>	<b>Effect on the organic matter content in soil on various time scales</b>		
	short	medium	long
Optimisation of crop rotation	++	++	++
Leaving crop residues behind	++	++	++
Supply of manure and compost	+++	++	+

Prevention of fallowing	+	++	+++
Lowering intensity and frequency of soil tillage	+	++	+++

*Table 1. Promising measures for the increase of organic matter content and their most important characteristics in the short (0 - 5 years), medium (5 – 10 years) and long (> 10 years) term.*

## 6. Costs and benefits

The additional costs of spreading garden- and food waste (VGF) or compost on the field instead of using animal slurry (which also contains organic matter) have been estimated by the Louis Bolk Institute to be between €150 (VGF) and €500 (compost) per ha per year ([De Wit, 2013](#)). The costs of green-manure crops amount to €130-€276 per ha per year including costs of sowing ([Kennisakker, 2017](#)). These cost key indicators may differ per type of soil, cultivation type (crop) and ways of utilisation.

Societal and economic benefits from increasing quantities of organic matter will be obtained by lowering irrigation costs, additional yield from the crop, carbon sequestration and effects on the water quality. The ‘state of the art’ on the agro-hydrological insights about this, has been described by [Groenendijk et al. \(2017\)](#). No economic interpretation is available of this research, thereby preventing drawing of proper conclusions about which conditions would justify, on a farming business scale, costs of application of green-manure plants, VGF or compost in order to raise organic matter contents.

## 7. Preconditions and promising locations

Organic matter is important for moisture supply, carrying capacity, water infiltration, nutrients storage, soil flora and fauna. This fact sheet specifically describes the role of organic matter on water management while in the fact sheet “[Bodem als Buffer](#)” the other positive effects of this measure are treated.

The direct effect of an increase of the organic matter content on water availability is especially relevant in a group of sandy soils with an organic matter percentage lower

than 3%. This means for the Netherlands in practice, that especially the green and yellow coloured sandy soils in figure 1, i.e. Mid- and East-Brabant, the Achterhoek and Twente, are promising locations.

The increase of soil organic matter content needs attention also in the long run, while it constitutes an investment with risks to be included in the decision-making of a regional water manager, who is busy realising a climate-proof water system.

## 8. Governance

The societal benefits of measures to increase the organic matter content on plot level, are difficult to express in terms of money or as a net revenue (costs-benefits). Yet it is clear that additional costs are involved, when the farmer wants to introduce such measures in his enterprise. An authority (water board, province) may consider to establish financial arrangements when additional costs do not just serve a private interest (more produce). Subsidies or interest-free loans can (partly) compensate for measures that lead to carbon sequestration or flooding prevention. Examples can be found in the Delta Freshwater programme, the Deltaplan Agrarian Water Management and POP3, as is summarised in the [Deltaprogramma Zoetwater 2017](#).

Water Boards in the Netherlands are more and more interested in and are contributing to measures, aimed at raising soil organic matter contents, like the [Lumbricus programme](#) and the Landbouwportaal Noord-Holland ([Waterschap Hollands Noorderkwartier](#), 2019). Much debate among board members still lingers, however, on who should bear the costs of soil-related measures on a plot level: the water manager or the agrarian entrepreneur.

## 9. Experiences from the Field

There is much direct or indirect attention, from research and other initiatives, for the buffering role of the soil, that retains water. An (incomplete) overview is given in paragraph 12. These initiatives are contributing to the realisation of the current [National Water Plan 2016-2021](#).

## 10. Knowledge Gaps

- Which cumulative effects play a role causing that measures taken, do not immediately result in a desired new equilibrium in the soil?
- How can the effect be quantified of an increase of organic matter content on soil functions like nutrients availability, trafficability, carbon sequestration, resilience against diseases and pests and crop production?
- What is the effect of increasing organic matter content on water availability at a plot-, farm- and river basin-level (scaling-up)?
- What are the economic and societal costs and benefits of measures aimed at increasing organic matter content?
- What should be the role division between water managers and farmers in stimulating process to actually embed measures for increasing organic matter?
- What is the influence of organic matter on soil biology? Effects to consider are permeability, stability, carrying capacity and water availability.

## 11. Sources & links

- [Akker, J.J.H. van den, F. de Vries, G.D. Vermeulen, M.J.D. Hack-ten Broeke en T. Schouten. 2012. Risico op ondergrondverdichting in het landelijk gebied in kaart. \(in Dutch\) Wageningen, Alterra, Alterra-Report 2409.](#)
- Berg, F. van den, A. Tiktak, T. Hoogland, A. Poot, J.J.T.I. Boesten, A.M.A. van der Linden, J.W. Pol. 2017. An improved soil organic matter map for GeoPEARL\_NL : Model description of version 4.4.4 and consequence for the Dutch decision tree on leaching to groundwater. Wageningen. [Wageningen Environmental Research report 2816.](#)
- Eurofins. 2017. Kwaliteit organische stof inzichtelijk gemaakt.[http://eurofins-agro.com/nl-nl/kwaliteit\\_organische\\_stof\\_inzichtelijk](http://eurofins-agro.com/nl-nl/kwaliteit_organische_stof_inzichtelijk)
- Faber, F., H. Wösten, G. Bakker, J. Bokhorst, E. Hummelink, I. Laros, . of fear Brink. Deru. and . of Eekeren. 2011. of grasslands in the. on https what see at others the sods of grass. Wageningen, Alterra-Rapport\_Alterra\_LBI 22 augustus. [ISSN 1566-7197. 116 p.](#)
- [Groenendijk., Schipper., van den Akker, J., Hendriks, R., Heinen, M. 2017. Effects improvement soil quality and water quality: Goede Soil for sustainable. Report Wageningen Environmental Research 2811.](#)
- [Kennisakker, 2017.](#)
- Klimaatberaad. 2018. [Ontwerp van het Klimaatakkoord](#). E. Nijpels (Ed.), (pp. 222). the Hague: Social Economic Board.

- Wageningen UR. 2012. [De Kwantitatieve Informatie Akkerbouw en Vollegrondsgroenteteelt 2012](#) (KWIN-AGV 2012) (pp. 203). Lelystad: Praktijkonderzoek Plant en Omgeving (Wageningen UR).
- [Reubens B., D., D., Ruysschaert G. 2010. Bodemkwaliteit en landbouw: a litterature review. Activiteit 1 van het Interregproject BodemBreed. Instituut voor Landbouw- en Visserijonderzoek \(ILVO\), Merelbeke-Lemberge, België. 203 p.](#)
- Rietberg, P.I., B. A. Visser. Kuikman. 2013. Manual good carbon management (in Dutch). Louis Bolk Institute, Driebergen, the Netherlands. Report 2013-002 LbP, pp. 29 ([http://www.louisboltk.org/nl/publicaties?ht\\_search=1&ht\\_also\\_above5years=1#books\\_and\\_reports](http://www.louisboltk.org/nl/publicaties?ht_search=1&ht_also_above5years=1#books_and_reports))
- [Rutgers M, Schouten T, Bloem J, Buis E, Dimmers W, van Eekeren N, de Goede RGM, Jagers op Akkerhuis GAJM, Keidel H, Korthals G, Mulder C, Wattel-Koekkoek EJW. 2015. Een indicatorsysteem voor ecosysteemdiensten van de bodem: Life support functions revisited. Report 2014-0145](#)
- Schouten. 2018. [Agriculture, nature and food: valuable and connected - the Netherlands as frontrunner in a circular agriculture. \(in Dutch pp. 37\)](#). Den Haag: Ministerie van Landbouw, Natuur en Voedselkwaliteit.
- [SmartSoil project. 2015. Sustainable farm management aimed at reducing threats to soils under climate change. European Union's Seventh Framework Programme, grant agreement no. 289694.](#)
- [Schipper et al. 2015. Goede grond voor een duurzaam watersysteem Verdere verkenningen in de relatie tussen agrarisch bodembeheer, bodemkwaliteit en waterhuishouding. Amersfoort, STOWA-Rapport 2015-19.](#)
- [Smit, A., Van Beek, C. L., & Hoogland, T. 2007. Risicogebieden voor organische stof - Ontwerp van een methodologie voor het aanwijzen van 'risk areas' t.b.v. de EU Kaderrichtlijn bodem \(pp. 51\). Wageningen: Wageningen UR.](#)
- [STOWA. 2015. Goede grond voor een duurzaam watersysteem', fase 1, Verdere verkenningen in de relatie tussen agrarisch bodembeheer, bodemkwaliteit en waterhuishouding. Authors: Piet Groenendijk, Peter Schipper \(Alterra\), Gijs Janssen en Joachim Rozemeijer \(Deltares\), Nick van Eekeren en Marleen Zanen \(Louis Bolk Instituut\) en Bjartur Swart\(ECS\).](#)
- TCB. 2016. [Advies Toestand en dynamiek van organische stof in Nederlandse landbouwbodems](#). Technische Commissie Bodem (J. v. Wensem (Ed.), rapport TCB A110, pp. 25. The Hague.

- [Van Eekeren, N., G.J. van der Burgt, B. Philipsen, H. van Schooten, M. de Haan. 2011. Effect op organische stof en kosten/baten, vruchtwisseling van gras en maïs, V-focus april 2011, pp. 24-26.](#)
- [Van Eekeren, N., Zaneveld-Reijnders, J. 2011. Bewust herstel van de natuurlijke buffercapaciteit van de bodem - Inhoudelijk rapportage 2010. Louis-Bolk Instituut & ZLTO.](#)
- Velthof, G.L., F.H. Kistenkas, P. Groenendijk, E.M.P.M. van Boekel en O. Oenema. 2018. [Wettelijk instrumentarium voor landbouwmaatregelen om waterkwaliteit te verbeteren. Realisatie van nutriëntendoelstellingen uit de Kaderrichtlijn Water.](#) Wageningen, WOT Nature & Environment, WUR. WOT-report 129. 118 pages.
- Water Board Hollands Noorderkwartier. 2019.  
<https://www.landbouwportaalnoordholland.nl/>
- [Williams et al., 2017. Organic Farming and Soil Physical Properties: An Assessment after 40 Years. Agron. J. 109: 600–609.](#)
- Wösten, J.H.M., G.J. Veerman, W.J.M. de Groot en J. Stolte. 2001. Waterretentie- en doorlatendheidskarakteristieken van boven- en ondergronden in Nederland: the Staringreeks. Renewed edition 2001 Alterra, Research Instituut voor de Groene Ruimte, Wageningen, [Alterra-rapport 153. 86 p.](#)
- Wit, J., de. 2013. [Bedrijfseconomische effecten van verhoging van het bodemorganische stofgehalte:](#) compostgebruik in de akkerbouw (pp. 17): Louis Bolk Institute.

*This factsheet is composed by Wageningen Environmental Research, February 2019*

**Authors:** Henk Wösten and Piet Groenendijk (Wageningen Environmental Research)

**Editor:** Jeroen Veraart (Wageningen Environmental Research)

**Translator:** Lidy van der Lugt (Wageningen Environmental Research)

## 12. Overview current initiatives and research

<b>Research project</b>	<b>Research location</b>	<b>Links/documents</b>
Soil Health Index	Wageningen Environmental Research	<a href="http://www.soilhealthindex.nl/">http://www.soilhealthindex.nl/</a>
SoilCare for profitable and sustainable crop production in Europe	Wageningen Environmental Research	<a href="https://www.soilcare-project.eu/">https://www.soilcare-project.eu/</a>
Recare: Preventing and remediating degradation of soil in Europe through land care	Wageningen Environmental Research	<a href="http://www.recare-project.eu/">http://www.recare-project.eu/</a>
Gezonde toplaag Gelderland (Louis Bolk Instituut, doorlopend)	Gelderland	<a href="http://www.louisbolt.org/nl/landbouw/bodembeheer/gezonde-toplaag-gelderland">http://www.louisbolt.org/nl/landbouw/bodembeheer/gezonde-toplaag-gelderland</a>
Goede grond voor een duurzaam watersysteem (2011-2015)	Nederland	<a href="http://www.deltaproof.stowa.nl/upload/publicatie2014/STOWA%202015%2019%20WEB%20DEF.pdf">http://www.deltaproof.stowa.nl/upload/publicatie2014/STOWA%202015%2019%20WEB%20DEF.pdf</a>
Gouden Gronden (continuous)	Noord Nederland	<a href="https://www.noorderzijlvest.nl/ons-werk/projecten/projecten-(lopend)/programma-gouden/">https://www.noorderzijlvest.nl/ons-werk/projecten/projecten-(lopend)/programma-gouden/</a>
Bodem kwaliteit op zandgrond (Vredepeel) (continuous)	PPO-locatie Vredepeel	<a href="https://www.wur.nl/nl/Onderzoek-Resultaten/Onderzoeksprojecten-LNV/Expertisegebieden/kennisonline/Bodemkwaliteit-op-zand-1.htm">https://www.wur.nl/nl/Onderzoek-Resultaten/Onderzoeksprojecten-LNV/Expertisegebieden/kennisonline/Bodemkwaliteit-op-zand-1.htm</a>
Duurzaam bodemgebruik veenweide (Veenweide Innovatie Centrum; Zegveld)	Veenweidegebieden (Utrecht)	<a href="https://www.veenweiden.nl/">https://www.veenweiden.nl/</a>
Deltaplan hoge zandgronden	Brabant/Waterschap Aa en Maas	<a href="https://www.brabant.nl/Dossiers/Dossiers-op-thema/Water/Bescherming-tegen-water/Rivieren-Maas-en-Merwedes/Deltaplan-Hoge-Zandgronden">https://www.brabant.nl/Dossiers/Dossiers-op-thema/Water/Bescherming-tegen-water/Rivieren-Maas-en-Merwedes/Deltaplan-Hoge-Zandgronden</a>
Kijk eens wat vaker onder de graszode	Vallei & Veluwe	<a href="http://www.grondwatercollectief.nl/portfolio-items/kijk-eens-wat-vaker-onder-de-graszode-meerwaarde-voor-boeren-en-waterbeheerders/">http://www.grondwatercollectief.nl/portfolio-items/kijk-eens-wat-vaker-onder-de-graszode-meerwaarde-voor-boeren-en-waterbeheerders/</a>

Bufferboeren – Samenwerken aan een betere waterbeschikbaarheid van de bodem	Bernheze (Noord-Brabant)	<a href="https://www.verantwoordeveehouderij.nl/show/Bufferboeren-Samen-werken-aan-een-betere-waterbeschikbaarheid-van-de-bodem.htm">https://www.verantwoordeveehouderij.nl/show/Bufferboeren-Samen-werken-aan-een-betere-waterbeschikbaarheid-van-de-bodem.htm</a>
Klimaatadaptief Waterbeheer: Wat biedt de bodem?	Stowa en SKB	Projects Information. <a href="http://edepot.wur.nl/221771">http://edepot.wur.nl/221771</a>
Gezond Zand	ROM3D	<a href="http://www.rom3d.nl/portfolio/gezond-zand/">http://www.rom3d.nl/portfolio/gezond-zand/</a>
Vruchtbare Kringloop	Rijn en IJssel	<a href="https://www.vruchtbarekringloop.nl/">https://www.vruchtbarekringloop.nl/</a>
Lumbricus	Aa en Maas/Vechtstromen	<a href="https://www.programmalumbricus.nl/">https://www.programmalumbricus.nl/</a>
Salland Waterproof	Waterschap Groot Salland	<a href="http://koolecommunicatie.nl/evaluatie-proces-salland-waterproof/">http://koolecommunicatie.nl/evaluatie-proces-salland-waterproof/</a>
Nationaal Kennis- en innovatieprogramma Water en Klimaat	Nederland	<a href="https://waterenklimaat.nl/nieuws/">https://waterenklimaat.nl/nieuws/</a>
Landbouw op Peil	Nederland	<a href="https://landbouwoppeil.nl/">https://landbouwoppeil.nl/</a>
Kenniscentrum Akkerbouw	Nederland	<a href="https://bo-akkerbouw.nl/">https://bo-akkerbouw.nl/</a>
Mijn bodemconditie	Nederland	<a href="http://mijnbodemconditie.nl/">http://mijnbodemconditie.nl/</a>

### 13. Disclaimer

De in deze publicatie gepresenteerde kennis en diagnosemethoden zijn gebaseerd op de meest recente inzichten in het vakgebied. Desalniettemin moeten bij toepassing ervan de resultaten te allen tijde kritisch worden beschouwd. De auteur(s) en STOWA kunnen niet aansprakelijk worden gesteld voor eventuele schade die ontstaat door toepassing van het gedachtegoed uit deze publicatie.

Knowledge and methods presented in this publication are based on the most recent insights in the discipline. Notwithstanding application of the results should be critically evaluated. The author(s) and STOWA cannot be taken responsible for possible damage caused by applying the results from this publication.